arXiv:1303.5350v1 [math.CO] 21 Mar 2013

PATTERN RECOGNITION ON ORIENTED MATROIDS: CRITICAL COMMITTEES AND DISTANCE SIGNALS

ANDREY O. MATVEEV

Abstract. If $\mathfrak{V}(\boldsymbol{R})$ is the vertex sequence of a symmetric cycle $\boldsymbol{R}$ in the tope graph of a simple acyclic oriented matroid $\mathcal{M}$ on a t-element ground set, then the set $\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})$ of minimal elements in the subposet $\mathfrak{V}(\boldsymbol{R})$ of the tope poset of $\mathcal{M}$, based at the positive tope, is a critical committee for $\mathcal{M}$ that votes for the base tope. We consider the sequence $\boldsymbol{z}_{\boldsymbol{R}} := (\rho(R) :\ R \in \mathfrak{V}(\boldsymbol{R}))$ of poset ranks of the elements from the vertex sequence of $\boldsymbol{R}$ as a fragment of a signal with period $2t$ and relate the number of members of the committee $\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})$ to the magnitudes of $\lfloor \frac{t}{2} \rfloor$ components, with odd indices, of the discrete Fourier transform of the distance vector $\boldsymbol{z}_{\boldsymbol{R}}$.

Contents

1. Introduction 1
2. Critical Committees and Distance Signals 2
References 6

1. Introduction

If $\mathfrak{V}$ is the vertex sequence of some path in a connected graph, then the corresponding sequence $(d(w, v) :\ v \in \mathfrak{V})$ of distances between its vertices and a distinguished vertex w of the graph can be extended in a natural way to a periodic signal. In this note, which is a companion to [3], we consider such distance signals associated with symmetric cycles in the tope graph of an oriented matroid. See [1] on oriented matroids. When the vertex sequence of a symmetric cycle is regarded as a subposet of the tope poset and the distinguished vertex of the tope graph is the base tope of the tope poset, a few basic observations (see, e.g., [2, Chapter 2],[5, Chapters 1÷6]) concerning the discrete Fourier transform (DFT) allow us to express the number of minimal elements of the vertex sequence of the symmetric cycle via the magnitudes of components of the DFT of the distance signal. In the case of an acyclic oriented matroid on a t-element ground set, with the distinguished positive tope, we thus relate the number of members of a

---

Key words and phrases. Discrete Fourier transform, critical committee, oriented matroid, tope, tope graph, tope poset.

2010 *Mathematics Subject Classification*: 52C40, 65T50.

critical committee to the magnitudes of $\lfloor\frac{t}{2}\rfloor$ components, with odd indices, of the DFT of the distance signal.

2. CRITICAL COMMITTEES AND DISTANCE SIGNALS

Let $\mathcal{A} := (E_t, \mathcal{T})$ be a simple acyclic oriented matroid on the ground set $E_t := \{1, \ldots, t\}$, with set of topes $\mathcal{T}$ whose components $-$ and $+$ are replaced by the real numbers -1 and 1, respectively. Let $\boldsymbol{R} := (R^0, R^1, \ldots, R^{2t-1}, R^0)$ be a symmetric cycle in the tope graph of $\mathcal{A}$, that is, $R^{k+t} = -R^k$, $0 \leq k \leq t-1$. Denote by $\mathcal{T}(\mathcal{L}(\mathcal{A}), \mathrm{T}^{(+)})$ the tope poset of $\mathcal{A}$ based at the positive tope $\mathrm{T}^{(+)}$. The set $\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})$ of minimal elements of the subposet $\mathfrak{V}(\boldsymbol{R}) := (R^0, R^1, \ldots, R^{2t-1}) \subset \mathcal{T}(\mathcal{L}(\mathcal{A}), \mathrm{T}^{(+)})$ is a *critical tope committee for* $\mathcal{A}$: it is the inclusion-minimal subset $\mathcal{K}^* \subset \mathfrak{V}(\boldsymbol{R})$ such that $\sum_{T \in \mathcal{K}^*} T = \mathrm{T}^{(+)}$. Let $\rho(T)$ denote the poset rank of a tope $T \in \mathcal{T}(\mathcal{L}(\mathcal{A}), \mathrm{T}^{(+)})$. The sequence

$$z_{\boldsymbol{R}} := \bigl(z_{\boldsymbol{R}}(0) := \rho(R^0), z_{\boldsymbol{R}}(1) := \rho(R^1), \ldots, z_{\boldsymbol{R}}(2t-1) := \rho(R^{2t-1})\bigr)$$

determines the *distance signal* $z_{\boldsymbol{R}} : \mathbb{Z} \to \{0\} \dot\cup E_t$ *of* $\boldsymbol{R}$, with period $2t$:

$$z_{\boldsymbol{R}}(j + 2t) = z_{\boldsymbol{R}}(j) , \quad j \in \mathbb{Z} .$$

Let $\ell^2(\mathbb{Z}_{2t})$ denote the $2t$-dimensional complex coordinate space; the elements of $\ell^2(\mathbb{Z}_{2t})$ are supposed to be row vectors whose components are indexed from 0 to $2t-1$. We consider the *distance vector* $z_{\boldsymbol{R}}$ *of* $\boldsymbol{R}$ as an element of the space $\ell^2(\mathbb{Z}_{2t})$.

Let $\boldsymbol{\iota}$ denote the vector $(1, 1, \ldots, 1) \in \ell^2(\mathbb{Z}_{2t})$. Since $z_{\boldsymbol{R}}(k) + z_{\boldsymbol{R}}(k+t) = t$, $0 \leq k \leq t-1$, we have

$$z_{\boldsymbol{R}} \cdot \boldsymbol{\iota}^\top = t^2 .$$

If $R \in \mathfrak{V}(\boldsymbol{R})$ is a vertex of the symmetric cycle $\boldsymbol{R}$ then we denote by $\mathcal{N}(R)$ the neighborhood of R in the cycle $\boldsymbol{R}$; $\mathbf{I}$ and $\mathbf{C}$ denote the $2t \times 2t$ identity matrix and basic circulant permutation matrix, respectively, with the rows and columns indexed from 0 to $2t-1$.

- On the one hand, $|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = t - \frac{1}{4} \sum_{\substack{\{R', R''\} := \mathcal{N}(R): \\ R \in \mathfrak{V}(\boldsymbol{R})}} |\rho(R'') - \rho(R')|$, and we have

$$\begin{aligned} |\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| &= t - \frac{1}{8} \sum_{\substack{\{R', R''\} := \mathcal{N}(R): \\ R \in \mathfrak{V}(\boldsymbol{R})}} \bigl(\rho(R'') - \rho(R')\bigr)^2 \\ &= t - \frac{1}{4} \sum_{\substack{\{R', R''\} := \mathcal{N}(R): \\ R \in \mathfrak{V}(\boldsymbol{R})}} \bigl(\rho(R)^2 - \rho(R')\rho(R'')\bigr) ; \end{aligned}$$

in other words, $|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = t - \frac{1}{4} z_{\boldsymbol{R}} \cdot (\mathbf{I} - \mathbf{C})(\mathbf{I} + \mathbf{C}) \cdot z_{\boldsymbol{R}}{}^\top$, or

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = t - \frac{1}{8} z_{\boldsymbol{R}} \cdot (2\mathbf{I} - \mathbf{C}^{-2} - \mathbf{C}^2) \cdot z_{\boldsymbol{R}}{}^\top . \tag{2.1}$$

The first row of the symmetric circulant matrix $2\mathbf{I}-\mathbf{C}^{-2}-\mathbf{C}^{2}$ is the vector $\mathbf{b} := (2,0,-1,0,\ldots,0,-1,0)$; the components of the DFT of $\mathbf{b}$ are

$$\hat{\mathrm{b}}(k) := \sum_{n=0}^{2t-1} \mathrm{b}(n)\mathsf{e}^{-\pi\imath kn/t} = 4\sin^2\tfrac{\pi k}{t}\ , \quad 0 \le k \le 2t-1\ .$$

Denote by $\mathbf{W}$ the $2t\times 2t$ Fourier matrix; its (m,n)-th entries are $\mathsf{e}^{-\pi\imath mn/t}$, $0\le m,n\le 2t-1$. The DFT and inverse DFT of the distance vector $\boldsymbol{z}_{\boldsymbol{R}}$ are the vectors $\hat{\boldsymbol{z}}_{\boldsymbol{R}} := \boldsymbol{z}_{\boldsymbol{R}}\mathbf{W}$ and $\check{\boldsymbol{z}}_{\boldsymbol{R}} := \boldsymbol{z}_{\boldsymbol{R}}\mathbf{W}^{-1}$, respectively; thus,

$$\begin{aligned}\hat{z}_{\boldsymbol{R}}(k) :&= \sum_{j=0}^{2t-1} z_{\boldsymbol{R}}(j)\mathsf{e}^{-\pi\imath kj/t}\\ &= \sum_{j=0}^{t-1}\bigl(z_{\boldsymbol{R}}(j)\mathsf{e}^{-\pi\imath kj/t} + (t-z_{\boldsymbol{R}}(j))\mathsf{e}^{-\pi\imath k(j+t)/t}\bigr)\ , \quad 0\le k\le 2t-1\ ,\end{aligned}$$

that is,

$$\hat{z}_{\boldsymbol{R}}(k) = \begin{cases} t^2\ , & k=0\ ,\\ 0\ , & k \text{ even}, \ k\neq 0\ ,\\ 2\bigl(-t(1-\mathsf{e}^{-\pi\imath k/t})^{-1} + \sum_{j=0}^{t-1} z_{\boldsymbol{R}}(j)\mathsf{e}^{-\pi\imath kj/t}\bigr)\ , & k \text{ odd}\ ;\end{cases}$$

in particular, if t is odd then $\hat{z}_{\boldsymbol{R}}(t) = -t + 2\sum_{j=0}^{t-1}(-1)^j z_{\boldsymbol{R}}(j)$.

We have

$$2\mathbf{I}-\mathbf{C}^{-2}-\mathbf{C}^{2} = \mathbf{W}^{-1}\cdot 4\,\mathrm{diag}(0,\sin^2\tfrac{\pi}{t},\ldots,\sin^2\tfrac{\pi k}{t},\ldots,\sin^2\tfrac{\pi(2t-1)}{t})\cdot\mathbf{W}\ ,$$

and Eq. (2.1) implies that

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = t - \tfrac{1}{2}\check{\boldsymbol{z}}_{\boldsymbol{R}}\cdot\mathrm{diag}(0,\sin^2\tfrac{\pi}{t},\ldots,\sin^2\tfrac{\pi k}{t},\ldots,\sin^2\tfrac{\pi(2t-1)}{t})\cdot\hat{\boldsymbol{z}}_{\boldsymbol{R}}^{\top}\ . \tag{2.2}$$

Denote by $\overline{\hat{\boldsymbol{z}}_{\boldsymbol{R}}}$ the vector composed of the complex conjugates of components of $\hat{\boldsymbol{z}}_{\boldsymbol{R}}$. Since $\check{\boldsymbol{z}}_{\boldsymbol{R}} = \frac{1}{2t}\overline{\hat{\boldsymbol{z}}_{\boldsymbol{R}}}$, it follows from Eq. (2.2) that

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = t - \tfrac{1}{4t}\overline{\hat{\boldsymbol{z}}_{\boldsymbol{R}}}\cdot\mathrm{diag}(0,\sin^2\tfrac{\pi}{t},\ldots,\sin^2\tfrac{\pi k}{t},\ldots,\sin^2\tfrac{\pi(2t-1)}{t})\cdot\hat{\boldsymbol{z}}_{\boldsymbol{R}}^{\top}\ .$$

Using Plancherel's formula, we reformulate this observation:

$$\begin{aligned}|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| &= t - \frac{1}{4t}\sum_{k=0}^{2t-1}|\hat{z}_{\boldsymbol{R}}(k)|^2\cdot\sin^2\tfrac{\pi k}{t} \qquad (2.3)\\ &= t - \frac{1}{2}\|\boldsymbol{z}_{\boldsymbol{R}}\|^2 + \frac{1}{4t}\sum_{k=0}^{2t-1}|\hat{z}_{\boldsymbol{R}}(k)|^2\cdot\cos^2\tfrac{\pi k}{t}\ .\end{aligned}$$

Since $\hat{z}_{\boldsymbol{R}}(k) = \overline{\hat{z}_{\boldsymbol{R}}(2t-k)}$ and the magnitudes $|\hat{z}_{\boldsymbol{R}}(k)|$ do not depend on circular translation: $|(\boldsymbol{z}_{\boldsymbol{R}}\mathbf{C}^{j})\hat{\ }(k)| = |\hat{z}_{\boldsymbol{R}}(k)|$, $1\le k\le 2t-1$, for any integer j, we restate Eq. (2.3) in the following way:

Proposition 2.1. (i) *For any distance vector* $\boldsymbol{z}_{\boldsymbol{R}}$ *of the symmetric cycle* $\boldsymbol{R}$*, we have*

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = t - \frac{1}{2t} \sum_{\substack{1 \leq k \leq t-1,\\ k \text{ odd}}} |\hat{z_{\boldsymbol{R}}}(k)|^2 \cdot \sin^2 \tfrac{\pi k}{t} \ . \tag{2.4}$$

(ii) *Let* $\mathcal{M} := (E_t, \mathcal{T})$ *be a simple oriented matroid (it has no loops, parallel or* antiparallel *elements), and* $\mathfrak{V}(\boldsymbol{R}) := (R^0, R^1, \ldots, R^{2t-1})$ *the vertex sequence of a symmetric cycle* $\boldsymbol{R}$ *in the tope graph of* $\mathcal{M}$*. Given a tope* T *of* $\mathcal{M}$*, denote by* $\boldsymbol{z}_{T,\boldsymbol{R}}$ *the* distance vector of $\boldsymbol{R}$ with respect to the tope T*, that is, the sequence* $(d(T,R) :\ R \in \mathfrak{V}(\boldsymbol{R}))$ *of graph distances between* T *and the vertices of* $\boldsymbol{R}$*. For the inclusion-minimal subset* $\boldsymbol{Q}(T,\boldsymbol{R}) \subset \mathfrak{V}(\boldsymbol{R})$ *such that* $T = \sum_{Q \in \boldsymbol{Q}(T,\boldsymbol{R})} Q$*, we have*

$$|\boldsymbol{Q}(T,\boldsymbol{R})| = t - \frac{1}{2t} \sum_{\substack{1 \leq k \leq t-1,\\ k \text{ odd}}} |\hat{z_{T,\boldsymbol{R}}}(k)|^2 \cdot \sin^2 \tfrac{\pi k}{t} \ . \tag{2.5}$$

The vertex sequence $\mathfrak{V}(\boldsymbol{R})$ is a maximal positive basis of $\mathbb{R}^t$; the existence of the set $\boldsymbol{Q}(T,\boldsymbol{R})$, of odd cardinality, of linearly independent elements of $\mathbb{R}^t$, mentioned in Proposition 2.1(ii), is guaranteed by [3, Corollary 2.2].

Following [4, Lecture 6], define the *distance enumerator*, with respect to a tope $B \in \mathcal{T}$, of the set $\mathcal{T}$, as the polynomial $D_{B,\mathcal{T}}(\mathrm{x}) := \sum_{T \in \mathcal{T}} \mathrm{x}^{d(B,T)}$. In an analogous manner, define the distance enumerator of the vertex set of the symmetric cycle $\boldsymbol{R}$ as the summand $D_{B,\mathfrak{V}(\boldsymbol{R})}(\mathrm{x}) := \sum_{T \in \mathfrak{V}(\boldsymbol{R})} \mathrm{x}^{d(B,T)}$ of $D_{B,\mathcal{T}}(\mathrm{x})$.

Note that for any topes T' and T'' of $\mathcal{M}$ the graph distance between them is $d(T',T'') = t - \frac{1}{4}\|T'' + T'\|^2 = \frac{1}{4}\|T'' - T'\|^2 = \frac{1}{2}\big(t - \langle T'', T' \rangle\big)$. Thus, if $\mathcal{T}^{\bullet}$ is a halfspace of $\mathcal{M}$, and if $\mathfrak{W} \subset \mathfrak{V}(\boldsymbol{R})$ is the vertex sequence of a $(t-1)$-path in the cycle $\boldsymbol{R}$, then $D_{B,\mathcal{T}}(\mathrm{x}) = \mathrm{x}^{t/2} \sum_{T \in \mathcal{T}^{\bullet}} (\mathrm{x}^{-\langle B,T \rangle/2} + \mathrm{x}^{\langle B,T \rangle/2})$ and $D_{B,\mathfrak{V}(\boldsymbol{R})}(\mathrm{x}) = \mathrm{x}^{t/2} \sum_{T \in \mathfrak{W}} (\mathrm{x}^{-\langle B,T \rangle/2} + \mathrm{x}^{\langle B,T \rangle/2})$.

Associate with a tope $T \in \mathcal{T}$ the $2t$-dimensional row vector $\boldsymbol{q}(T,\boldsymbol{R})$ defined by $q_j(T,\boldsymbol{R}) := 1$ if $R^j \in \boldsymbol{Q}(T,\boldsymbol{R})$, and $q_j(T,\boldsymbol{R}) := 0$ otherwise. Let $\mathbf{G}(\boldsymbol{R})$ denote the Gram matrix of the sequence $\mathfrak{V}(\boldsymbol{R})$. For topes T' and T'' of $\mathcal{M}$ we have $d(T',T'') = \frac{1}{2}\big(t - \boldsymbol{q}(T'',\boldsymbol{R})\mathbf{G}(\boldsymbol{R})\boldsymbol{q}(T',\boldsymbol{R})^{\top}\big)$.

• On the other hand,

$$\begin{aligned} |\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| &= \frac{1}{8} \sum_{\substack{\{R',R''\} := \mathcal{N}(R):\\ R \in \mathfrak{V}(\boldsymbol{R})}} \big(\rho(R') + \rho(R'') - 2\rho(R)\big)^2 \\ &= \frac{1}{4} \sum_{\substack{\{R',R''\} := \mathcal{N}(R):\\ R \in \mathfrak{V}(\boldsymbol{R})}} \big(3\rho(R)^2 - 2\rho(R)(\rho(R') + \rho(R'')) + \rho(R')\rho(R'')\big) \ , \end{aligned}$$

that is, $|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{1}{4}\boldsymbol{z}_{\boldsymbol{R}} \cdot (\mathbf{I} - \mathbf{C})(3\mathbf{I} - \mathbf{C}) \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top}$, or

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{1}{8}\boldsymbol{z}_{\boldsymbol{R}} \cdot (6\mathbf{I} - 4\mathbf{C}^{-1} - 4\mathbf{C} + \mathbf{C}^{-2} + \mathbf{C}^2) \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top} \ . \tag{2.6}$$

Eqs. (2.1) and (2.6) imply that $|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{t}{2} + \frac{1}{4}\boldsymbol{z}_{\boldsymbol{R}} \cdot (\mathbf{I}-\mathbf{C})^2 \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top}$ $= \frac{3t}{4} - \frac{1}{4}\boldsymbol{z}_{\boldsymbol{R}} \cdot (\mathbf{I}-\mathbf{C})\mathbf{C} \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top}$; in other words, we have

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{t}{2} + \frac{1}{8}\boldsymbol{z}_{\boldsymbol{R}} \cdot (2\mathbf{I} - 2\mathbf{C}^{-1} - 2\mathbf{C} + \mathbf{C}^{-2} + \mathbf{C}^2) \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top} \tag{2.7}$$

and

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{3t}{4} - \frac{1}{8}\boldsymbol{z}_{\boldsymbol{R}} \cdot (\mathbf{C}^{-1} + \mathbf{C} - \mathbf{C}^{-2} - \mathbf{C}^2) \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top} \;; \tag{2.8}$$

note also that

$$\boldsymbol{z}_{\boldsymbol{R}} \cdot (\mathbf{I}-\mathbf{C}) \cdot {\boldsymbol{z}_{\boldsymbol{R}}}^{\top} = t \; .$$

We derive from Eqs. (2.6), (2.7) and (2.8) the relations

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{1}{4t}\sum_{k=0}^{2t-1} |\hat{z}_{\boldsymbol{R}}(k)|^2 \cdot (\cos^2 \tfrac{\pi k}{t} - 2\cos\tfrac{\pi k}{t} + 1) \; ,$$

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{t}{2} + \frac{1}{4t}\sum_{k=0}^{2t-1} |\hat{z}_{\boldsymbol{R}}(k)|^2 \cdot (\cos^2 \tfrac{\pi k}{t} - \cos\tfrac{\pi k}{t})$$

and

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R})| = \frac{3t}{4} + \frac{1}{8t}\sum_{k=0}^{2t-1} |\hat{z}_{\boldsymbol{R}}(k)|^2 \cdot (2\cos^2 \tfrac{\pi k}{t} - \cos\tfrac{\pi k}{t} - 1)$$

that are equivalent to Eq. (2.3).

• Let $\boldsymbol{R}'$ and $\boldsymbol{R}''$ be two symmetric cycles in the tope graph of the oriented matroid $\mathcal{A}$, and $\boldsymbol{z}_{\boldsymbol{R}'}$ and $\boldsymbol{z}_{\boldsymbol{R}''}$ their distance vectors. Consider the vectors $\boldsymbol{e} := \boldsymbol{z}_{\boldsymbol{R}''} - \boldsymbol{z}_{\boldsymbol{R}'}$ and $\boldsymbol{m} := \boldsymbol{z}_{\boldsymbol{R}''} + \boldsymbol{z}_{\boldsymbol{R}'}$. Note that the components of their DFTs are:

$$\hat{e}(k) = \begin{cases} 0 \; , & k \text{ even} \; , \\ 2\sum_{j=0}^{t-1} e(j)\mathsf{e}^{-\pi\imath kj/t} \; , & k \text{ odd} \; , \end{cases}$$

and

$$\hat{m}(k) = \begin{cases} 2t^2 \; , & k = 0 \; , \\ 0 \; , & k \text{ even}, \; k \neq 0 \; , \\ 2\bigl(-2t(1-\mathsf{e}^{-\pi\imath k/t})^{-1} + \sum_{j=0}^{t-1} m(j)\mathsf{e}^{-\pi\imath kj/t}\bigr) \; , & k \text{ odd} \; ; \end{cases}$$

thus, if t is odd then $\hat{e}(t) = 2\sum_{j=0}^{t-1}(-1)^j e(j)$ and $\hat{m}(t) = 2\bigl(-t + \sum_{j=0}^{t-1}(-1)^j m(j)\bigr)$.

It follows from Proposition 2.1(i) that

$$|\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R}')| + |\,\mathbf{min}\,\mathfrak{V}(\boldsymbol{R}'')| = 2t - \frac{1}{4t}\sum_{\substack{1\leq k\leq t-1, \\ k \text{ odd}}} \bigl(|\hat{e}(k)|^2 + |\hat{m}(k)|^2\bigr) \cdot \sin^2 \tfrac{\pi k}{t} \; .$$

References

[1] A. Björner, M. Las Vergnas, B. Sturmfels, N. White and G.M. Ziegler, *Oriented Matroids*, Encyclopedia of Mathematics, **46**, Cambridge University Press, Cambridge, 1993. Second edition 1999.

[2] M.W. Frazier, *An Introduction to Wavelets through Linear Algebra*, Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1999.

[3] A.O. Matveev, *Pattern Recognition on Oriented Matroids: Topes and Critical Committees*, `arXiv:1011.6082`.

[4] R.P. Stanley, *An Introduction to Hyperplane Arrangements*, Geometric Combinatorics, 389–496, IAS/Park City Math. Ser., 13, Amer. Math. Soc., Providence, RI, 2007.

[5] M.W. Wong, *Discrete Fourier Analysis*, Pseudo-Differential Operators. Theory and Applications, 5. Birkhäuser/Springer Basel AG, Basel, 2011.

E-mail address: `andrey.o.matveev@gmail.com`